
Patient Advisory Information Labels











A-2

THIS MEDICINE MAY IMPAIR YOUR ABILITY

A-16

May Cause DROWSINESS I-106



TAKE THIS MEDICINE ON AN EMPTY STOMACH, 1 HOUR BEFORE OR 2 HOURS AFTER A MEAL, UNLESS OTHERWISE DIRECTED BY YOUR DOCTOR.

A-3



TAKE WITH FOOD OR MILK I-108




DO NOT TAKE ANTACIDS,

DAIRY PRODUCTS OF IRON PREPARATIONS

2 HOURS BEFORE OR


2 HOURS AFTER THIS MEDICINE.















DRINK PLENTY OF WATER WHILE TAKING THIS MEDICINE. A-5



I-111

effects to FDA at 1-800-FDA-1088. I-134

THIS MEDICINE MAY DISCOLOR YOUR URINE or FECES.



This prescription CANNOT be REFILLED without a WRITTEN PRESCRIPTION from your DOCTOR.

THIS PRESCRIPTION HAS **BEEN COMPOUNDED BY PHARMACY**

A-6





I-113

40156









A-8



I-114

40157









IMPORTANT: AVOID PROLONGED OR AND/OR ARTIFICIAL SUNLIGHT
WHILE TAKING THIS MEDICINE.


FOR THE EYE Use This Medicine Exactly as Directed by Your Doctor. A-48



I-119



A-11







I-102 A-12

